

Convergence of the Ishikawa Iteration Process for Nonexpansive Mappings

Lei Deng

*Department of Mathematics, Southwest China Normal University, Beibei,
 Chongqing 630715, People's Republic of China*

E

y Elsevier - Publisher Connector

Received March 1, 1995

Let D be a subset of a normed space X and $T: D \rightarrow X$ be a nonexpansive mapping. Given a sequence $\{x_n\}$ in D and two real sequences $\{t_n\}$ and $\{s_n\}$ satisfying

- (i) $0 \leq t_n \leq t < 1$ and $\sum_{n=1}^{\infty} t_n = \infty$,
- (ii) $0 \leq s_n \leq 1$ and $\sum_{n=1}^{\infty} s_n < \infty$,
- (iii) $x_{n+1} = t_n T(s_n T x_n + (1 - s_n)x_n) + (1 - t_n)x_n$, $n = 1, 2, 3, \dots$,

we prove that if $\{x_n\}$ is bounded, then $\lim_{n \rightarrow \infty} \|T x_n - x_n\| = 0$. The conditions on D , X , and T are shown which guarantee the weak and strong convergence of the Ishikawa iteration process to a fixed point of T . © 1996 Academic Press, Inc.

1. INTRODUCTION

Let D be a subset of a normed space X and $T: D \rightarrow X$ be nonexpansive (this is, $\|Tx - Ty\| \leq \|x - y\|$ for all x, y in D). In 1974, Ishikawa [5] introduced a new iteration process as

$$x_{n+1} = t_n T(s_n T x_n + (1 - s_n)x_n) + (1 - t_n)x_n, \quad n = 1, 2, 3, \dots,$$

where $\{t_n\}$ and $\{s_n\}$ are sequences in $[0, 1]$ satisfying certain restrictions. The Mann iteration process is a special case of Ishikawa where $s_n = 0$ for all $n \geq 1$ [1, 4, 7].

In 1976, Ishikawa [6] proved the following theorem without any assumption on convexity of the Banach space.

THEOREM A [6]. *Let D be a subset of a normed space X and $T: D \rightarrow X$ be a nonexpansive mapping. Given a sequence $\{x_n\}$ in D and a sequence $\{t_n\}$*

of real numbers satisfying

(i) $0 \leq t_n \leq t < 1$ and $\sum_{n=1}^{\infty} t_n = \infty$,

(ii) $x_{n+1} = (1 - t_n)x_n + t_nTx_n$, for $n = 1, 2, 3, \dots$,

if $\{x_n\}$ is bounded, then $\|Tx_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

In this paper, we first extend Theorem A to the Ishikawa iteration process. Then we show the weak and strong convergence of the Ishikawa iteration process in a Banach space. Our results generalize the results of Emmauele [2] and Ishikawa [6].

2. LEMMAS

LEMMA 1 [11]. Suppose that $\{a_n\}$ and $\{b_n\}$ are two sequences of nonnegative numbers such that $a_{n+1} \leq a_n + b_n$ for all $n \geq 1$. If $\sum_{n=1}^{\infty} b_n$ converges, then $\lim_{n \rightarrow \infty} a_n$ exists.

LEMMA 2. Suppose that $\{a_n\}$ and $\{b_n\}$ are two sequences of a normed space X . If there is a sequence $\{t_n\}$ of real numbers satisfying

(i) $0 \leq t_n \leq t < 1$ and $\sum_{n=1}^{\infty} t_n = \infty$,

(ii) $a_{n+1} = (1 - t_n)a_n + t_nb_n$ for all $n \geq 1$

(iii) $\lim_{n \rightarrow \infty} \|a_n\| = d$,

(iv) $\limsup_{n \rightarrow \infty} \|b_n\| \leq d$ and $\{\sum_{i=1}^n t_i b_i\}$ is bounded,

then $d = 0$.

Proof. Suppose that $d > 0$ and it follows from (iv) that $\sum_{i=n}^{n+m-1} t_i b_i$ is bounded for all n and m . Let

$$M = \sup \left\{ \left\| \sum_{i=n}^{n+m-1} t_i b_i \right\| : n, m = 1, 2, 3, \dots \right\}.$$

Choose a number N such that

$$N > \max \left\{ \frac{2M}{d}, 1 \right\}.$$

We can choose a positive ε such that

$$1 - 2\varepsilon \exp \left(\frac{N+1}{1-t} \right) > \frac{1}{2}.$$

It follows from (i) that there exists a natural k such that

$$N < \sum_{i=1}^k t_i \leq N + 1.$$

Since $\lim_{n \rightarrow \infty} \|a_n\| = d$, $\limsup_{n \rightarrow \infty} \|b_n\| \leq d$ and ε independent of n , without loss of generality we may assume that for all $n \geq 1$,

$$d(1 - \varepsilon) < \|a_n\| < d(1 + \varepsilon) \quad \text{and} \quad \|b_n\| < d(1 + \varepsilon).$$

Setting $T = \sum_{i=1}^k t_i$, $S = \prod_{i=1}^k s_i$ and $s_n = 1 - t_n$ for all $n \geq 1$, from (ii), we obtain

$$a_{k+1} = s_1 s_2 \dots s_k a_1 + t_1 s_2 s_3 \dots s_k b_1 + \dots + t_{k-1} s_k b_{k-1} + t_k b_k,$$

$$a_{k+1} \in B := \text{co}\{a_1, b_1, b_2, \dots, b_k\}.$$

Let $x = T^{-1} \sum_{i=1}^k t_i b_i$ and

$$y = S(1 - S)^{-1} \{a_1 + t_1(s_1^{-1} - T^{-1})b_1 \\ + t_2(s_1^{-1}s_2^{-1} - T^{-1})b_2 + \dots + t_k(S^{-1} - T^{-1})b_k\}.$$

Then it is clear that $x, y \in B$ and $a_{k+1} = Sx + (1 - S)y$. Therefore

$$d(1 - \varepsilon) < \|a_{k+1}\| \leq S\|x\| + (1 - S)\|y\| \\ \leq S\|x\| + (1 - S)d(1 + \varepsilon).$$

Hence, we have

$$\begin{aligned} \|x\| &> d(1 - S^{-1}(2 - S)\varepsilon) > d(1 - 2\varepsilon S^{-1}) \\ &= d\left(1 - 2\varepsilon \prod_{i=1}^k (1 - t_i)^{-1}\right) \\ &= d\left[1 - 2\varepsilon \exp\left(\sum_{i=1}^k \log\left(1 + \frac{t_i}{1 - t_i}\right)\right)\right] \\ &\geq d\left(1 - 2\varepsilon \exp\left(\sum_{i=1}^k \frac{t_i}{1 - t_i}\right)\right) \\ &\geq d\left(1 - 2\varepsilon \exp\left(\frac{T}{1 - t}\right)\right) \\ &\geq d\left(1 - 2\varepsilon \exp\left(\frac{N + 1}{1 - t}\right)\right) > \frac{d}{2}, \end{aligned}$$

since $\log(1 + u) \leq u$ for $-1 < u < \infty$.

On the other hand, we have

$$\|x\| = T^{-1} \left\| \sum_{i=1}^k t_i b_i \right\| \leq T^{-1} M \leq \frac{d}{2M} M = \frac{d}{2},$$

arriving at a contradiction. This completes the proof.

LEMMA 3 [11]. *Let D be a subset of a normed space X and $T: D \rightarrow X$ be a nonexpansive mapping. Given a sequence $\{x_n\}$ in D satisfying*

$$x_{n+1} = t_n T(s_n T x_n + (1 - s_n) x_n) + (1 - t_n) x_n, \quad n = 1, 2, 3, \dots,$$

where $0 \leq t_n, s_n \leq 1$ for all $n \geq 1$, then

$$\|x_{n+1} - p\| \leq \|x_n - p\| \quad \text{for all } n \geq 1 \text{ and all } p \in F(T),$$

where $F(T)$ denotes the set of fixed points of T .

3. MAIN RESULTS

THEOREM 1. *Let D be a subset of a normed space X and $T: D \rightarrow X$ be a nonexpansive mapping. Given a sequence $\{x_n\}$ in D and two real sequences $\{t_n\}$ and $\{s_n\}$ satisfying*

$$(i) \quad 0 \leq t_n \leq t < 1 \text{ and } \sum_{n=1}^{\infty} t_n = \infty,$$

$$(ii) \quad 0 \leq s_n \leq 1 \text{ and } \sum_{n=1}^{\infty} s_n < \infty,$$

$$(iii) \quad x_{n+1} = t_n T(s_n T x_n + (1 - s_n) x_n) + (1 - t_n) x_n, \quad n = 1, 2, 3, \dots,$$

if $\{x_n\}$ is bounded, then $\|T x_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Set $y_n = s_n T x_n + (1 - s_n) x_n$, then $x_{n+1} = t_n T y_n + (1 - t_n) x_n$. By page 304 of [11], we have

$$\|T x_{n+1} - x_{n+1}\| \leq (1 + 2s_n(1 - t_n)) \|T x_n - x_n\|.$$

Setting $a_n = T x_n - x_n$, by (ii) and the fact that $\{\|T x_n - x_n\|\}$ is bounded, it follows from Lemma 1 that $\lim_{n \rightarrow \infty} \|a_n\| = d$.

We may assume that $t_n > 0$ for all $n \geq 1$, without loss of generality. Setting $b_n = t_n^{-1}(T x_{n+1} - T x_n) + T x_n - T y_n$, we have $a_{n+1} = (1 - t_n) a_n + t_n b_n$ and

$$\begin{aligned} \|b_n\| &\leq t_n^{-1} \|T x_{n+1} - T x_n\| + \|T x_n - T y_n\| \\ &\leq t_n^{-1} \|x_{n+1} - x_n\| + \|x_n - y_n\| \\ &= \|T y_n - x_n\| + s_n \|T x_n - x_n\| \\ &\leq \|T y_n - T x_n\| + \|T x_n - x_n\| + s_n \|T x_n - x_n\| \\ &\leq \|T x_n - x_n\| + 2s_n \|T x_n - x_n\|. \end{aligned}$$

By (ii), $\lim_{n \rightarrow \infty} s_n = 0$, we have

$$\limsup_{n \rightarrow \infty} \|b_n\| \leq d.$$

Finally, we have

$$\begin{aligned} \left\| \sum_{i=1}^n t_i b_i \right\| &= \left\| \sum_{i=1}^n (Tx_{i+1} - Tx_i + t_i(Tx_i - Ty_i)) \right\| \\ &= \left\| Tx_{n+1} - Tx_1 + \sum_{i=1}^n t_i(Tx_i - Ty_i) \right\| \\ &\leq \|Tx_{n+1} - Tx_1\| + \sum_{i=1}^n t_i \|Tx_i - Ty_i\| \\ &\leq \|x_{n+1} - x_1\| + \sum_{i=1}^n t_i s_i \|Tx_i - x_i\| \end{aligned}$$

which is bounded. Hence it follows from Lemma 2 that $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$, completing the proof.

Theorem 1 above generalizes Lemma 2 of Ishikawa [6].

Recall that a Banach space X is said to satisfy Opial's condition [8] if the condition $x_n \rightarrow x_0$ weakly implies

$$\limsup_{n \rightarrow \infty} \|x_n - x_0\| < \limsup_{n \rightarrow \infty} \|x_n - y\|$$

for all $y \neq x_0$.

THEOREM 2. *Let X be a Banach space which satisfies Opial's condition, D be weakly compact, and let T and $\{x_n\}$ be as in Theorem 1. Then $\{x_n\}$ converges weakly to a fixed point of T .*

Proof. Due to weak compactness of D , there exists $\{x_{n_k}\}$ of $\{x_n\}$ which converges weakly to a $p \in D$. With standard proof we show that $p = Tp$. We suppose that $\{x_n\}$ doesn't converge weakly to p ; then there are $\{x_{n_j}\}$ and $q \neq p$ such that $x_{n_j} \rightarrow q$ weakly and $q = Tq$. By Lemma 3 and Opial's condition of X , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - p\| &= \lim_{k \rightarrow \infty} \|x_{n_k} - p\| < \lim_{k \rightarrow \infty} \|x_{n_k} - q\| \\ &= \lim_{j \rightarrow \infty} \|x_{n_j} - q\| < \lim_{j \rightarrow \infty} \|x_{n_j} - p\| \\ &= \lim_{n \rightarrow \infty} \|x_n - p\|, \end{aligned}$$

a contradiction. This completes the proof.

Theorem 2 above generalizes a result of Emmanuele [2].

Let D be a subset of a Banach space X . A mapping $T: D \rightarrow X$ with a nonempty fixed points set $F(T)$ in D will be said to satisfy Condition A [10] if there is a nondecreasing function $f: [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$, $f(r) > 0$ for $r \in (0, \infty)$, such that $\|x - Tx\| \geq f(d(x, F(T)))$ for all $x \in D$, where $d(x, F(T)) = \inf\{\|x - z\|: z \in F(T)\}$.

The two following theorems generalize Theorems 1 and 2 of Ishikawa [6], respectively. Since a similar proof is in [6, 11], we omit their proof here.

THEOREM 3. *Let D be a closed subset of a Banach space X , and T be a nonexpansive mapping from D into a compact subset of X . If $\{x_n\}$ is as in Theorem 1, then $\{x_n\}$ converges to a fixed point of T .*

THEOREM 4. *Let X , D , and $\{x_n\}$ be as in Theorem 3. Let $T: D \rightarrow X$ be a nonexpansive mapping with a nonempty fixed points set $F(T)$ in D . If T satisfies Condition A, then $\{x_n\}$ converges to a member of $F(T)$.*

The following theorem generalizes Theorem 1.8 of Veeramani [12].

THEOREM 5. *Let D be a closed, bounded, convex subset of a Banach space X and $T: D \rightarrow D$ be a map satisfying*

- (i) $\|Tx - Ty\| \leq \|x - y\|$, for all $x, y \in D$;
- (ii) For some $\alpha > 0$,

$$\|Tx - Ty\| \leq \alpha(\|x - Tx\| + \|y - Ty\|), \quad \text{for all } x, y \in D.$$

If $\{x_n\}$ is as in Theorem 1, then $\{x_n\}$ converges to the unique fixed point of T .

Proof. By Gillespie and Williams [3], it follows that T has a unique fixed point p in D . From (ii), we have

$$\|Tx_n - p\| = \|Tx_n - Tp\| \leq \alpha\|x_n - Tx_n\|.$$

Hence, by Theorem 1, we obtain

$$\|Tx_n - p\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

It follows that

$$\|x_n - p\| \leq \|x_n - Tx_n\| + \|Tx_n - p\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This completes the proof.

The following theorem extends Theorem 2 of Reich [9] in uniformly convex Banach spaces.

THEOREM 6. *Suppose that X is a uniformly convex Banach space with a Frechet differentiable norm, C a nonempty closed convex subset of X ,*

$T: C \rightarrow C$ a nonexpansive mapping with a fixed point, and $\{x_n\}$ is as in Theorem 1. Then $\{x_n\}$ converges weakly to a fixed point of T .

Proof. Let $\omega_w(x_n)$ be the weak limit ω -set of $\{x_n\}$, i.e., the set $\{x \in X: x = \text{weak-lim}_{k \rightarrow \infty} x_{n_k} \text{ for some } n_k \uparrow \infty\}$. By Theorem 1 and Browder's demiclosedness principle, $\omega_w(x_n)$ is contained in $F(T)$, the fixed point set of T .

The remainder of the proof is similar to that of Theorem 1 of [11], so the details are omitted.

ACKNOWLEDGMENT

The author is grateful to the referee for his careful reading and valuable suggestions which appended Theorem 6.

REFERENCES

1. W. G. Dotson, Jr., On the Main iterative process, *Trans. Amer. Math. Soc.* **149** (1970), 65–73.
2. G. Emmanuele, Convergence of the Mann-Ishikawa iterative process for nonexpansive mappings, *Nonlinear Anal.* **6** (1982), 1135–1141.
3. A. A. Gillespie and B. B. Williams, Some theorems on fixed points in Lipschitz and Kannan type mappings, *J. Math. Anal. Appl.* **74** (1980), 382–387.
4. C. W. Groetsch, A note on segmenting Mann iterates, *J. Math. Anal. Appl.* **40** (1972), 369–372.
5. S. Ishikawa, Fixed points by a new iteration method, *Proc. Amer. Math. Soc.* **44** (1974), 147–150.
6. S. Ishikawa, Fixed points and iterations of a nonexpansive mapping in a Banach space, *Proc. Amer. Math. Soc.* **59** (1976), 65–71.
7. W. R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* **4** (1953), 506–510.
8. Z. Opial, Weak convergence of the sequence of successive approximations for nonexpansive mappings, *Bull. Amer. Math. Soc.* **73** (1967), 595–597.
9. S. Reich, Weak convergence theorem for nonexpansive mappings in Banach spaces, *J. Math. Anal. Appl.* **67** (1979), 274–276.
10. H. F. Senter and W. G. Dotson, Jr., Approximating fixed points of nonexpansive mappings, *Proc. Amer. Math. Soc.* **44** (1974), 375–380.
11. K. K. Tan and H. K. Xu, Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process, *J. Math. Anal. Appl.* **178** (1993), 301–308.
12. P. Veeramani, On some fixed point theorems on uniformly convex Banach spaces, *J. Math. Anal. Appl.* **167** (1992), 160–166.